ASB Standard 039, First Edition 20212023

Standard for Internal Validation of Human Short Tandem Repeat Profiling on Capillary Electrophoresis Platforms



This document is copyrighted © by the AAFS Standards Board, LLC. <u>20212023</u> All rights are reserved. 410 North 21st Street, Colorado Springs, CO 80904, www.aafs.org/academy-standards-board.

Standard for Internal Validation of Human Short Tandem Repeat Profiling on Capillary Electrophoresis Platforms

ASB Approved Xxxxx 20212023

ANSI Approved Xxxxxx 20212023



410 North 21st Street Colorado Springs, CO 80904

This document may be downloaded from: www.aafs.org/academy-standards-board

This document is provided by the AAFS Academy Standards Board. Users are permitted to print and download the document and extracts from the document for personal use, however the following actions are prohibited under copyright:

- modifying this document or its related graphics in any way;
- using any illustrations or any graphics separately from any accompanying text; and,
- failing to include an acknowledgment alongside the copied material noting the AAFS Academy Standards Board as the copyright holder and publisher.

Users may not reproduce, duplicate, copy, sell, resell, or exploit for any commercial purposes this document or any portion of it. Users may create a hyperlink to <u>www.aafs.org/academy-standards-board</u> to allow persons to download their individual free copy of this document. The hyperlink must not portray AAFS, the AAFS Standards Board, this document, our agents, associates and affiliates in an offensive manner, or be misleading or false. ASB trademarks may not be used as part of a link without written permission from ASB.

The AAFS Standards Board retains the sole right to submit this document to any other forum for any purpose.

Certain commercial entities, equipment or materials may be identified in this document to describe a procedure or concept adequately. Such identification is not intended to imply recommendations or endorsement by the AAFS or the AAFS Standards Board, nor is it intended to imply that the entities, materials, or equipment are necessarily the best available for the purpose.

Proper citation of ASB documents includes the designation, title, edition, and year of publication.

This document is copyrighted[®] by the AAFS Standards Board, LLC. 20212023 All rights are reserved. 410 North 21st Street, Colorado Springs, CO 80904, www.aafs.org/academy-standards-board.

Foreword

This document outlines the requirements for internal validation of short tandem repeat DNA profiling using STR kits on capillary electrophoresis platforms utilized in forensic laboratories.

The validation of capillary electrophoresis is one part of the process of generating a DNA result and there are steps prior to and after that need to be considered.

The American Academy of Forensic Sciences established the Academy Standards Board (ASB) in 2015 with a vision of safeguarding Justice, Integrity and Fairness through Consensus Based American National Standards. To that end, the ASB develops consensus based forensic standards within a framework accredited by the American National Standards Institute (ANSI), and provides training to support those standards. ASB values integrity, scientific rigor, openness, due process, collaboration, excellence, diversity and inclusion. ASB is dedicated to developing and making freely accessible the highest quality documentary forensic science consensus Standards, Guidelines, Best Practices, and Technical Reports in a wide range of forensic science disciplines as a service to forensic practitioners and the legal system.

This document was revised, prepared, and finalized as a standard by the DNA Consensus Body of the AAFS Standards Board. The draft of this standard was developed by the Human Forensic Biology Subcommittee of the Organization of Scientific Area Committees (OSAC) for Forensic Science.

Questions, comments, and suggestions for the improvement of this document can be sent to AAFS-ASB Secretariat, <u>asb@aafs.org</u> or 401 N 21st Street, Colorado Springs, CO 80904.

All hyperlinks and web addresses shown in this document are current as of the publication date of this standard.

ASB procedures are publicly available, free of cost, at <u>www.aafs.org/academy-standards-board</u>.

Keywords: internal validation, DNA, short tandem repeat profiling, capillary electrophoresis.

Table of Contents (to be finalized prior to publication)

l

| 1 | Scope | |
|------------------------------------|-----------------------|--|
| 2 | Normative References | |
| 3 | Terms and Definitions | |
| 4 | Requirements | |
| 5 | Conformance | |
| Annex A (informative) Bibliography | | |
| | | |

Standard for Internal Validation of Human Short Tandem Repeat Profiling on Capillary Electrophoresis Platforms

1 Scope

This document details requirements for performing an internal validation of a human short tandem repeat (STR) multiplex kit using capillary electrophoresis (CE].) in forensic laboratories.

2 Normative References

There are no normative reference documents. Annex A, Bibliography, contains informative references.

The following normative reference is indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments applies.

ANSI/ASB Standard 038, *Standard for Internal Validation of Forensic DNA Analysis Methods*, First Edition, 2020^a.

3 Terms and Definitions

For purposes of this document, the following definitions apply.

3.1

analytical threshold

The minimum height requirement at and above which detected peaks on a STR DNA profile electropherogram can be reliably distinguished from instrument background noise; peaks above this threshold are generally not considered noise and are either artifacts or true alleles.

3.2

artifact

A non-allelic product of the amplification process (e.g., stutter, non-templated nucleotide addition, or other non-specific product), an anomaly of the detection process (e.g., single or multichannel voltage spikes or "pull-up"), or a by-product of primer synthesis (e.g., "dye blob") that may be observed on an electropherogram.

3.3

contamination

Exogenous DNA or other biological material in a DNA sample, PCR reaction, or item of evidence; the exogenous DNA or biological material could be present before the sample is collected, or introduced during collection or testing of the sample.

^a Available from: https://www.aafs.org/academy-standards-board

3.4

drop-in

Allelic peak(s) in an electropherogram that are not reproducible across multiple independent amplification events.

3.5

drop-out

Failure of an otherwise amplifiable allele to produce a signal above analytical threshold because the allele was not present in sufficient quantity in the aliquot that underwent PCR amplification.

3.6

interpretation

The process of evaluating DNA data for purposes including, but not limited to, defining assumptions related to mixtures and single source profiles, distinguishing between alleles and artifacts, assessing the possibility of degradation, inhibition, and stochastic effects, and determining whether the data are suitable for comparison.

<u>3.63.7</u>

peak height ratio

The relative ratio of two peaks at a given locus in a diploid heterozygous single-source sample.

<u>3.73.8</u>

precision

The degree of mutual agreement among a series of individual measurements, values and/or results^b.

<u>3.83.9</u>

repeatability studies

Studies to evaluate the degree of variability in multiple measurements (e.g_{-u} replicate samples) under constant conditions, such as studies by the same operator using the same equipment in the same laboratory within short intervals of time.

<u>3.93.10</u>

reproducibility studies

Studies to evaluate the degree of variability in multiple measurements (e.g., replicate samples) under varying conditions, such as studies using the same methods with different operators or different equipment.

<u>3.103.11</u>

sensitivity studies

Studies performed during developmental and/or internal validation of DNA or other test methods designed to define the lower and upper limits/bounds of an assay to accurately detect an analyte.

<u>3.113.12</u>

stochastic threshold

The peak height value in a DNA profile above which it is reasonable to assume that, at a given locus, allelic drop-out of a sister allele in a heterozygous pair has not occurred in a single source DNA sample: due to the possibility of shared alleles in mixed samples, the presence of allele peaks above

<u>https://www.swgdam.org/#!publications/c1mix</u>

the stochastic threshold is no guarantee that allele dropout did not occur in mixed DNA sample profiles.

<u>3.12</u>3.13

stutter

An artifact of polymerase chain reaction (PCR) amplification typically observed one or more repeat units smaller or larger than a short tandem repeat (STR) allele in a DNA profile, may result from strand slippage during PCR amplification. A stutter peak is generally of lower relative fluorescence units (RFU) than the allele peak.

<u>3.133.14</u>

validation

The process of performing and evaluating a set of experiments that <u>establishestablishes</u> the efficacy, reliability, and limitations of a method, procedure or modification thereof; establishing recorded documentation that provides a high degree of assurance that a specific process will consistently produce an outcome meeting its predetermined specifications and quality attributes. May include developmental and/or internal validation.

4 Requirements

4.1 The internal validation of STR test kits using capillary electrophoresis should include the following studies and mayshall include the studies described in this section, in conjunction with ANSI/ASB Standard 038 and supported by relevant developmental validation studies. This work shall also be supplemented with published scientific literature or other appropriate scientific resources, where available. See additional materials available to include the ASB Best Practice Recommendation 129 (not yet published, also out for public comment) included in Annex A.

A single analysis software package (e.g., GeneMapper^{™c} or GeneMarker^{®d}), version and set of analysis parameters shall be used throughout the performance of studies described below. If changes are made to the analysis parameters, the impact of those changes shall be described and where needed the data reanalyzed.

4.1.1 The laboratory shall conduct an STR genotyping concordance study.

4.1.1.1 STR genotypes obtained from the new STR test kit shall be compared to genotypes obtained from the previous kit for all overlapping loci, if applicable.

4.1.1.2 This study shall include the evaluation of concordance of the new STR test kit utilizing an appropriate certified reference material for STR genotyping.

4.1.1.3 In the event that a discordant genotype is observed when comparing identical genetic markers, this discordant data shall be noted in the final validation document, and a potential explanation for the discordant data supported by data or other research studies shall be included in the final validation document.

<u>^c</u> This term is used as an example only, and does not constitute an endorsement of this product by the AAFS Standards Board.

^d This term is used as an example only, and does not constitute an endorsement of this product by the AAFS Standards Board.

4.1.2 The laboratory shall perform sensitivity studies that will include replicates for each set of assay conditions/parameters (i.e.g., PCR cycle number, injection time/, injection voltage, and PCR reaction volume, etc.) that the lablaboratory utilizes with the new STR kit.

4.1.2.1 This study shall include a range of DNA inputs anticipated to be amplified as well as lower and higher quantities to empirically define the upper and lower limits of DNA inputs to be used for testing.

4.1.2.2 DNA extracts spanning a range of DNA quality shall be included- (e.g., synthetically and/or naturally degraded as well as inhibited DNA templates of known types expected to be encountered by the testing laboratory).

NOTE Optimal input ranges <u>for single source DNA samples</u> can be determined using the studies described in 4.1.2.1 and 4.1.2.2.

4.1.3 The laboratory shall determine the analytical threshold for each dye channel of the new STR test kit using a range of sample types and DNA input quantities across multiple analyses.

4.1.4 The laboratory shall characterize expected peak height ratio variation observed for each locus of the new STR test kit utilizing single source samples amplified over a range of DNA input amounts.

4.1.5 The laboratory shall characterize expected PCR stutter artifacts observed for each STR locus of the new STR test kit.

4.1.6 The laboratory shall determine stochastic threshold(s) for any manual binary interpretation method used to interpret the new STR test kit data.

4.1.7 The laboratory shall conduct a <u>studystudies</u> utilizing mixed DNA samples <u>usinghaving</u> a range of DNA ratios, DNA template input quantities, and <u>numbernumbers</u> of contributors <u>with</u> <u>varied degrees of allele sharing</u> expected to be interpreted by the testing laboratory.

The results of this studythese studies shall be used to establish define the limits of mixture interpretation and to establish the protocols for determiningassessing the number of contributors, to be used in the interpretation, the use of mixture ratios, and distinguishing major and minor contributors components of mixtures.

<u>NOTE</u> Depending on the interpretation method chosen, additional studies may need to be performed (e.g., specificity testing within probabilistic genotyping). Additional information on mixture studies and the development of interpretation protocols can be found in ASB Standard 020.

4.1.8 The laboratory shall demonstrate allelic sizing precision and calling accuracy of the new STR test kit and capillary electrophoresis instrument through repeatability and reproducibility studies.

4.1.9 The laboratory shall determine the susceptibility of the genotyping process to the introduction and detection of exogenous DNA by documenting allelic drop-in and contamination through the evaluation of controls (i.e., reagent blanks, negative and positive amplification controls) and samples with known genotypes. The laboratory shall document drop-in and contamination events and calculate drop-in rates in accordance with their ts quality system.

4.1.10 The validation study results shall be used to determine the standard operating procedure of the laboratory for this kit. Verification of the laboratory's newly established standard operating procedures using the new STR test kit shall be performed shall conduct studies utilizing known and casework-like samples with a range of sample types representative of those regularly analyzed expected to be encountered by the testing laboratory.

4.1.10.1 The laboratory shall compare the results to previous results of known and casework-like samples, where applicable.

4.1.10.2<u>4</u>**.1.9.1** This study shallshould minimally include processing a set of samples using all DNA extraction and quantification chemistries utilized by the laboratory to verify compatibility with the new STR test kit.

4.1.9.2 The results of these studies shall be used to evaluate the thresholds and other parameters that have been established during previous studies and where possible, address the limitations of the protocols to be included in the standard operating procedures for this kit.

5 Conformance

In order to demonstrate conformance with this standard, the laboratory shall have completed and reviewed the following:

<u>a) documentation of</u> all <u>applicableinternal</u> validation studies as listed in Section 4. The; <u>b) documented quality assurance parameters, interpretation protocols, and analytical procedures</u> <u>derived from internal</u> validation studies<u>; and</u>

<u>c) documented approval by the DNA Technical Leader or other appropriate personnel prior to implementation and communicated to all analysts.</u>

<u>All documentation and data from studies performed</u> shall be summarized and<u>made readily</u> available for review- (e.g., by auditors or inspectors, stakeholders who use reports generated by the DNA mixture test protocols and parameters).

Annex A (informative)

Bibliography

The following information provides a list of the literature resources that may assist the DNA technical leader in defining the breadth and scope of the validation materials that may be useful. This list is not meant to be all inclusive. The laboratory shall develop a list tailored to its specific needs. Updated references shall be added to the laboratory's list as new methods or technologies are incorporated into the laboratory's protocols.

- 1] ANSI/ASB Standard 018, Standard for Validation of Probabilistic Genotyping Systems, First Edition, 2020^e.
- 2] ANSI/ASB Standard 020, Standard for Validation Studies of DNA Mixtures, and Development and Verification of a Laboratory's Mixture Interpretation Protocol, First Edition 2018^a
- 3] ANSI/ASB Standard 038, Standard for Internal Validation of Forensic DNA Analysis Methods, First Edition 2020^d.
- 4] ANSI/ASB Standard 040, Standard for Forensic DNA Interpretation and Comparison Protocols, First Edition, 2019^d.
- 5] ASB Std 129 Reference to be added upon publication.
- 1]6] Bregu, J., D. Conklin, E. Coronado, M. Terrill, R.W. Cotton, and C.M. Grgicak, Analytical "Thresholds and Sensitivity: Establishing RFU Thresholds for Forensic DNA Analysis." *Journal* of Forensic Science, 2013, 58, pp. 120–129.
- 2]7] Butler, J.M. "Quality Assurance and Validation." *In: Advanced Topics in Forensic DNA Typing: Methodology.* Elsevier, 2011.
- <u>3]8]</u> ENSFI. *Recommended Minimum Criteria for the Validation of Various Aspects of the DNA* Profiling Process. 2010.^f.
- 4<u>]9]</u>Gilder, J. R., T.E. Doom, K. Inman, and D.E. Krane, "Run-Specific Limits of Detection and Quantitation for STR-based DNA Testing." *Journal of Forensic Sciences*, 2007, 52, pp. 97–101. doi:10.1111/j.1556-4029.2006.00318.x
- 5]10] FBI, Quality Assurance Standards for DNA Databasing Laboratories. Effective July 1, 2020g.
- 6]11] FBI, Quality Assurance Standards for Forensic DNA Testing Laboratories. Effective July 1, 2020^h.

<u>^g Available from: https://docs.wixstatic.com/ugd/4344b0_809d01b3e9f9451cb9edd9a85f2c2e5b.pdf</u>

^e Available from: https://www.aafs.org/academy-standards-board

f Available from: http://enfsi.eu/wp-content/uploads/2016/09/minimum validation guidelines in dna profiling - v2010 0.pdf

hAvailable from: https://docs.wixstatic.com/ugd/4344b0 6782472e073442ec877085584aaffa36.pdf

- 7]12] SWGDAM. SWGDAM Guidelines for STR Enhanced Detection Methodsⁱ.
- 8]13] SWGDAM. SWGDAM Interpretation Guidelines for Autosomal STR Typing by Forensic DNA Testing Laboratoriesⁱ.
- <u>9]14]</u> SWGDAM. SWGDAM Interpretation Guidelines for Y-Chromosome Testing^k.
- 10]15]_SWGDAM. SWGDAM Validation Guidelines for Forensic DNA Analysis Methods¹.

ⁱAvailable from: http://media.wix.com/ugd/4344b0_29feed748e3742a5a7112467cccec8dd.pdf ^jAvailable from: https://docs.wixstatic.com/ugd/4344b0_50e2749756a242528e6285a5bb478f4c.pdf ^kAvailable from: http://media.wix.com/ugd/4344b0_da25419ba2dd4363bc4e5e8fe7025882.pdf ^lAvailable from: https://docs.wixstatic.com/ugd/4344b0_813b241e8944497e99b9c45b163b76bd.pdf



Academy Standards Board 410 North 21st Street Colorado Springs, CO 80904

www.aafs.org/academy-standards-boardwww.asbstandardsboard.org