ASB Standard 147, First Edition 20222023

# Standard for Analyzing Skeletal Trauma in Forensic Anthropology



This document is copyrighted<sup>©</sup> by the AAFS Standards Board, LLC. <del>2022</del>2023 All rights are reserved. 410 North 21st Street, Colorado Springs, CO 80904, www.aafs.org/academy-standards-board.

# Standard for Analyzing Skeletal Trauma in Forensic Anthropology

ASB Approved Xxxxx 20222023

ANSI Approved Xxxxxx 20222023



410 North 21<sup>st</sup> Street Colorado Springs, CO 80904

This document may be downloaded from: www.aafs.org/academy-standards-board

This document is provided by the AAFS Academy Standards Board. Users are permitted to print and download the document and extracts from the document for personal use, however the following actions are prohibited under copyright:

- modifying this document or its related graphics in any way;
- using any illustrations or any graphics separately from any accompanying text; and,
- failing to include an acknowledgment alongside the copied material noting the AAFS Academy Standards Board as the copyright holder and publisher.

Users may not reproduce, duplicate, copy, sell, resell, or exploit for any commercial purposes this document or any portion of it. Users may create a hyperlink to <u>www.aafs.org/academy-standards-board</u> to allow persons to download their individual free copy of this document. The hyperlink must not portray AAFS, the AAFS Standards Board, this document, our agents, associates and affiliates in an offensive manner, or be misleading or false. ASB trademarks may not be used as part of a link without written permission from ASB.

The AAFS Standards Board retains the sole right to submit this document to any other forum for any purpose.

Certain commercial entities, equipment or materials may be identified in this document to describe a procedure or concept adequately. Such identification is not intended to imply recommendations or endorsement by the AAFS or the AAFS Standards Board, nor is it intended to imply that the entities, materials, or equipment are necessarily the best available for the purpose.

Proper citation of ASB documents includes the designation, title, edition, and year of publication.

This document is copyrighted<sup>©</sup> by the AAFS Standards Board, LLC. <del>20222023</del> All rights are reserved. 410 North 21st Street, Colorado Springs, CO 80904, www.aafs.org/academy-standards-board.

# Foreword

This standard was developed to provide guidance to practitioners for recognizing, describing, interpreting, and reporting trauma in skeletal material. Trauma analysis is a component of anthropologic examination needed to identify traumatic events occurring before and during death and differentiate from damage occurring after death. This document is intended to assist forensic anthropologists when assessing trauma from skeletal elements.

The American Academy of Forensic Sciences established the Academy Standards Board (ASB) in 2015 with a vision of safeguarding Justice, Integrity and Fairness through Consensus Based American National Standards. To that end, the ASB develops consensus based forensic standards within a framework accredited by the American National Standards Institute (ANSI), and provides training to support those standards. ASB values integrity, scientific rigor, openness, due process, collaboration, excellence, diversity and inclusion. ASB is dedicated to developing and making freely accessible the highest quality documentary forensic science consensus Standards, Guidelines, Best Practices, and Technical Reports in a wide range of forensic science disciplines as a service to forensic practitioners and the legal system.

This document was revised, prepared, and finalized as a standard by the Anthropology Consensus Body of the AAFS Standards Board. The draft of this standard was developed by the Anthropology Subcommittee of the Organization of Scientific Area Committees (OSAC) for Forensic Science.

Questions, comments, and suggestions for the improvement of this document can be sent to AAFS-ASB Secretariat, <u>asb@aafs.org</u> or 401 N 21st Street, Colorado Springs, CO 80904.

All hyperlinks and web addresses shown in this document are current as of the publication date of this standard.

ASB procedures are publicly available, free of cost, at <u>www.aafs.org/academy-standards-board</u>.

Keywords: trauma analysis; skeletal examination; trauma timing; trauma mechanism

# Table of Contents (to be updated when the document is finalized)

l

1	Scope
2	Normative References
3	Terms and Definitions
4.1	Requirements General Trauma Timing
	Trauma Mechanism
5	Considerations and Adjustments
6	Reporting

# Standard for Analyzing Skeletal Trauma in Forensic Anthropology

# 1 Scope

This standard provides requirements for documenting, describing, interpreting, and reporting skeletal trauma in forensic anthropology. It also provides requirements for the determination of trauma timing (i.e., antemortem, perimortem, or postmortem) and the identification of the mechanism that produced the trauma (i.e., projectile, sharp, blunt, or thermal trauma). This document does not address cause and manner of death.

# 2 Normative References

There are no normative reference documents.

# 3 Terms and Definitions

For purposes of this document, the following definitions apply.

## 3.1

# antemortem trauma

A skeletal defect that occurred before an individual's death and shows evidence of osteological reaction.

## 3.2

## blunt trauma

A skeletal defect produced by low-velocity impact from a blunt object (e.g., being struck by an object or concussive wave) or the low-velocity impact of a body with a blunt surface (e.g., motor vehicle accident or fall).

## 3.3

## delamination

A separation of cortical and cancellous bone.

## 3.4

## kerf

The notch or groove in bone, tooth, or cartilage made by an object that is edged, pointed, or beveled.

## 3.5

## perimortem trauma

A skeletal defect that occurred when bone is in a biomechanically fresh (visco-elastic) state with no evidence of an osteological reaction.

## 3.6

## plastic deformation

A permanent distortion of bone's shape without fracture caused by a force exceeding an element's elastic limit/elastic modulus/yield point.

## 3.7

## postmortem damage

A skeletal defect that occurred after a bone has lost its biomechanically fresh (visco-elastic) properties.

## 3.8

## projectile trauma

A skeletal defect often produced by high-velocity impact over a relatively small surface area, typically by projectiles from firearms, but can result from any small object impacting a bone at a high velocity.

## 3.9

## sharp trauma

A skeletal defect produced by an object that is edged, pointed, or beveled.

## 3.10

## skeletal trauma

A defect to osseous, dental, and/or cartilaginous tissue as a result of external forces.

## 3.11

# spatial distribution

The locations of defects (or damage/breakage) in relation to each other throughout the skeleton.

## 3.12

## thermal trauma

A skeletal defect produced by exposure to high temperature or direct contact with flame.

## 3.13

## trauma description

Reporting of the location, dimension(s), and other observed characteristics of the skeletal defect(s).

## 3.14

## trauma interpretation

Explanation regarding the mechanism, timing, direction of impact(s<del>), and/</del>or minimum number of impacts associated with skeletal defect(s) based onusing quantitative and/or qualitative observations.evidence

### 3.15

## trauma mechanism

The external factors that produce a skeletal defect.

## **4** Requirements

## 4.1 General

Skeletal material shall be assessed and evidence of trauma and damage shall be documented before and after processing. The cranium should be opened and a complete endocranial examination conducted. Ideally, the endocranial examination should occur after ectocranial trauma is documented and other cranial data are collected. SkeletalWhen possible, skeletal trauma examination shall be examined using scientifically validatedbased on gross, microscopic, radiographic, photographic, and/or other analyticalobservations and analyzed using peer reviewed published methods. Documentation of trauma shall be adequate to allow independent verification of work performed in the absence of the originally analyzed material.

Documentation of skeletal trauma shall include written descriptions as well as visual representations (i.e., photographs, diagrams/sketches, radiographs, casts, 3D scanning). Descriptions shall include the location and characteristics of the skeletal defect. Trauma location shall be documented using standard osteological/anatomical terms. When multiple defects are present, the spatial distribution shall be documented and described. Dimensions of skeletal defects resulting from trauma should be measured, when appropriate.

Foreign material (e.g., hair, fibers, bullet fragments, blade tip) associated with a skeletal defect shall be documented and described. These materials may be collected and submitted for further analysis by other forensic science providers. However, analysis of foreign material is beyond the scope of this document.

A distinction shall be drawn between trauma description and interpretation. At a minimum, a description of the defect(s) shall be given. Trauma interpretation shall be limited to when the evidence supports the findings.

# 4.2 Trauma Timing

## 4.2.1 General

Defect characteristics shall be used to assess when the skeletal element was traumatized or damaged. Timing of the defect shall be classified using terms such as antemortem, perimortem, or postmortem. When a distinction in timing cannot be made between antemortem, perimortem, and postmortem, the limitations shall be clearly documented and reported.

## 4.2.2 Antemortem Trauma

Antemortem trauma shall be identified based on the presence of a trauma-related osteological reaction. These reactions may include healed fractures or evidence of healing, the development of pseudarthrosis, degenerative joint disease, or infectious response related to a fracture, a dental fracture with worn or rounded fracture margin, or a surgically implanted device.

Practitioners shall not provide a trauma interpretation for antemortem trauma except in cases that show identifiable features and patterning or radiographic evidence of identifiable foreign bodies.

## 4.2.3 Perimortem Trauma

Perimortem trauma shall be identified based on the presence of biomechanical characteristics indicative of fresh bone or the incontrovertible association of the trauma with the incident that produced it (e.g., explosion, fire, plane crash, fall from a building). Presence of biomechanical characteristics shall include: lack of osteological reaction, presence of fresh bone fracture characteristics, and absence of dry bone fracture characteristics. Classification of trauma as perimortem is strengthened when the mechanism or indicators can be identified (e.g., hair

entrapped in bone with associated blunt trauma injuries, the presence of blood staining, or embedded debris from a fall from a building).

#### 4.2.4 Postmortem Damage

Postmortem damage shall be identified based on differentially stained or recently exposed surfaces, absence of healing, characteristics of the break lacking biomechanically fresh (visco-elastic) properties, and changes resulting from taphonomic events.

When defects are classified as postmortem, terms such as "damage" and "breakage" should be used. The terms "trauma" and "fracture" should be reserved for antemortem and perimortem trauma.

#### 4.3 Trauma Mechanism

### 4.3.1 General

Trauma mechanisms shall be based on defect shape and size, spatial distribution, and evidence of plastic deformation. Extrinsic, continuously variable and concurrent factors such as velocity and force may preclude the identification of a trauma mechanism. When the trauma mechanism cannot be identified, the defect shall be clearly described, documented, and reported without interpretation.

Fractured bones shall be re-approximated, and may be reconstructed, to clearly assess defect features and their spatial distribution. Fracture margins and fracture surfaces shall be examined prior to reconstruction and a reversible medium should be used. When possible, observed fracture patterns should be compared to reference material to aid in trauma mechanism classification.

No conclusions regarding defect sequencing shall be provided unless there are clear indications of the order of defects (e.g., a fracture terminating at a preexisting fracture).

### 4.3.2 Projectile Trauma

Observed features indicating projectile trauma may include: a projectile in association with the bone, entrance or exit defect characteristics (e.g<sub>7.1</sub> internal or external beveling), projectile residue, bullet wipe, remnants of the projectile, fracture pattern with minimal to no plastic deformation, and external beveling of concentric fractures in the cranial vault that indicates an internal to external force.

In the context of projectile trauma, plastic deformation may indicate a lower velocity projectile, a projectile slowed before impact, or other energy loss during flight.

When possible, entrance and exit defects shall include descriptions and photographs of internal and external beveling, defect measurement, anatomic location, associated fracture patterns (e.g., radiating and concentric fractures), and estimation of projectile path relative to anatomical position.

The presence of projectile residue (e.g., soot and other materials discharged from a firearm, staining from extended contact with metal, embedded metallic/radiodense artifacts) shall be documented. The term "bullet wipe" is preferred to "lead wipe," since the outer surface of a bullet may be composed of materials other than lead.

Practitioners shall not estimate or report bullet caliber. Estimates of muzzle to target distance shall be made with caution and supported by skeletal findings. Practitioners should describe the spread of shotgun pellets, presence of wadding, or shot cup, which may be informative of distance.

#### 4.3.3 Blunt Trauma

Observed features of blunt trauma may include: fracture patterns (e.g. wedge, buckle), plastic deformation, delamination, internal beveling of concentric fractures in the cranial vault indicative of external to internal force. Features of an impact site (i.e., depressed fractures, circumferential fractures, tool marks) shall be documented. Staining, which may result from blood and fat within crushed diploe, and other alterations which may represent impact sites shall also be documented.

The use of standard clinical terms (e.g., Parry, Colles) and orthopedic classification systems (e.g., Le Fort, Salter-Harris fractures) should be used with caution to avoid non-skeletal implications or interpretation.

An interpretation of the minimum number of impacts and direction of impact(s) should be documented, when possible.

#### 4.3.4 Sharp Trauma

Observed features of sharp trauma may include: straight-line incised defects, punctures or gouges, chop or hack marks (clefts), and kerfs. Defect features such as length, width, depth, and interstriation distance should be measured, and casts made when possible. Sharp trauma defects and their casts shall be examined grossly and microscopically.

Defect features may reflect class characteristics of the tool and shall be interpreted to identify tool type/class. Dismemberment should be considered when interpreting sharp force trauma. particularly if the extremities and/or head are affected.

Fractures can occur in association with sharp trauma and shall be documented, if present. Features common with pseudo-sharp trauma, such as scrapes, scores, and scratch marks, shall be documented. Tool marks in bone or cartilage shall not be identified as hesitation marks.

### 4.3.5 Thermal Trauma

Observed features of thermal trauma may include: color changes (e.g., yellow, black, white), delamination, burn pattern, shrinkage, charring, calcination, or heat-related fracture patterns. Thermal fracture location(s) and type(s) shall be documented. Features indicating that the bone was biomechanically dry during burning (e.g., absence of warping) shall be documented. When thermal trauma and other trauma (e.g., sharp, blunt, projectile) coexist, all trauma types shall be clearly delineated in the documentation. Aberrant patterns of burning (i.e., patterns that are inconsistent with those expected due to tissue shielding) may provide information about body positioning, presence of trauma (e.g., dismemberment), or postmortem interval (e.g., lack of pugilistic posture) and shall be documented.

Practitioners shall not estimate the temperature or duration of heat exposure based on thermal defects to bone.

## 5 Considerations and Adjustments

The term "perimortem" is used differently by pathologists and anthropologists. The timing of injury is less precise when evaluating bone than when evaluating soft tissue. Based upon the intrinsic properties of bone, the perimortem interval extends from days preceding death to days to weeks following the death event. When perimortem trauma is identified, a forensic anthropological definition of "perimortem" should be included in the report.

Practitioners may estimate the minimum number of traumatic events (e.g., blunt impacts, projectile entry defects, or sharp defects) observed skeletally, but shall not report a definitive maximum number of impacts, as skeletal trauma evidence may not reflect all impacts to the body.

The spatial distribution of fractures shall be considered to evaluate whether the fractures occurred from a single impact/event (e.g., a fall resulting in serial rib fractures).

Factors that may influence skeletal trauma (e.g., decedent age, sex, health status; bone and fracture type; fracture location) shall be considered in skeletal trauma interpretation. These factors are particularly important when assessing age of injury. For example, an adult healing phase assessment shall not be applied to an infant.

Pathological conditions, anomalies, or taphonomic changes may mimic skeletal trauma.

Practitioners shall not determine a "match" between a specific tool and a tool mark.

Forensic anthropologists shall not determine cause or manner of death.

Blasts/explosive events often cause blunt (including concussive) and projectile trauma to the body. When the trauma pattern and circumstantial information are is consistent with a blast event, the trauma mechanism should be classified as ""\_blast trauma." ".

## 6 Reporting

The written report shall include the methods used and descriptions of gross, microscopic, and radiological findings that are referenced in support of a trauma interpretation. Trauma location and spatial distribution as related to anatomical position shall be reported.

Interpretation of trauma and its timing may not be possible due to the nature of the trauma or the condition of the skeletal material; equivocal results shall be reported as such. Trauma interpretation shall be clearly identified in the report using terms such as 'indicative of' and 'consistent with' or by using a subheading titled 'Interpretation'.

When a suspect tool is submitted for analysis, similarities between the tool and defect may be reported; conclusions shall be reported in terms of an exclusion or failure to exclude.

Reporting shall avoid terminology that may be considered inflammatory, imply a particular outcome, or imply intent. Terms to avoid include but are not limited to: victim, weapon, violent, painful, suffer, and lethal.

When casts are made of traumatic defects, the method, materials, and any artifacts created, as well as the final disposition, shall be documented in the report or case file. When physical evidence is

collected from a traumatic defect (e.g., a projectile, blade tip), the disposition of the evidence shall be documented.

l

7



Academy Standards Board 410 North 21st Street Colorado Springs, CO 80904

www.aafs.org/academy-standards-board