



A131 Improved Isolation of DNA From Forensic Dental Specimens

Allison E. Reklaitis, MS*, and Chenique Sanders, BS*, Office of the Chief Medical Examiner, New York City Office, 421 East 26th Street, Department of Forensic Biology, New York, NY 10016; My K. Tran, DDS, and Amr M. Moursi, DDS, New York University College of Dentistry, Pediatrics Department, 345 East 24th Street, New York, NY 10016; and Sheila E. Dennis, MS, Office of the Chief Medical Examiner, New York City Office, Department of Forensic Biology, 421 East 26th Street, New York, NY 10016

After attending this presentation, attendees will know a different approach to the examination of forensic dental specimens and also the effect of specific environmental conditions on the ability to extract DNA from these specimens, specifically teeth.

This presentation will impact the forensic science community by giving those who work on forensic dental specimens and other specimens, that are compromised by the environmental conditions they are exposed to, another approach to examination and knowledge that the three specific environments that these teeth were exposed to do have an impact on the quality of DNA obtained.

The DNA Missing Persons Unit within the Department of Forensic Biology at the New York City Office of Chief Medical Examiner performs nuclear and mitochondrial DNA testing for the purposes of identification, re-association of body parts, and upload into the Combined DNA Index System (CODIS). While success rates for postmortem samples such as blood, tissue and bone were as expected, dental specimens failed to give consistent results. This may have been caused by exposure to detrimental environmental conditions. These environments may vary from bodies found in water, to remains buried in soil, or encased in cement. The goal of this study is to evaluate how exposure to the following conditions: water, soil, and, cement, affects the ability to extract nuclear DNA from dental samples; and determine what improvements can be made to the examination and extraction protocols.

One hundred-twenty adult teeth collected as dental waste from private dental offices were examined and categorized by a dentist according to tooth type (molars, incisors, premolars, and canines) and any abnormalities such as dental caries (cavities), open apices, calcification, and discoloration were noted. The teeth were incubated in soil, cement, water from the East River in New York City, or physiological saline from one to 48 weeks. Two specimens from each environmental condition had the pulp tissue extracted at Weeks one, two, three and four and every four weeks thereafter using a dental pulpectomy procedure. The pulpectomy involved removing tissue from the pulp chambers using nerve broaches utilized by dentists when performing a root canal. A phenol chloroform isoamyl alcohol (PCIA) extraction was performed on the pulp tissue. Each DNA sample was quantified, amplified with the Applied Biosystems (ABI) Identifiler[®] kit (28 cycles), and alleles detected using the ABI PRISM 3130xl Genetic Analyzer.

The current method used to extract DNA from teeth involved grinding the entire tooth using a freezer mill. This destroyed the morphology of the tooth and successful DNA results fluctuated and were unpredictable. Results from this study greatly improved examination procedures; the morphology of the tooth structure was maintained while examination efficiency increased by accessing the DNA in the tooth through pulpectomies.

By week 40, 30.8% generated full profiles and 16.7% generated high partial profiles all eligible for entry into CODIS. Overall by week 40, the number of identified loci decreased over time in most conditions but DNA extracted from dental specimens buried in soil showed the strongest reduction in identifiable loci.

Prior caries status of a tooth was not a factor in DNA extraction as those teeth yielded sufficient amounts of DNA. Conversely, teeth with open apices, that were calcified, or had no visible pulp tissue did not yield DNA or yield sufficient amounts of DNA for concentration or amplification.

Overall, the aims of this study thus far have been met. Improvements to the examination of dental specimens through pulpectomies require less examination and hands on time compared to the conventional freezer mill method. Pulpectomies preserve the morphology of the tooth and yield similar DNA results. The dental specimens exposed to cement, soil, water, and even physiological saline show degradation over time as evidenced by the decrease in DNA concentration and quality of DNA profiles obtained.

DNA, Extraction, Identification