



## Physical Anthropology Section - 2012

### H33 A New Method for Histological Age Estimation of the Femur

*Christian Crowder, PhD\*, and Victoria M. Dominguez, MA, Office of the Chief Medical Examiner, 520 First Avenue, New York, NY 10016*

After attending this presentation, attendees will understand the current issues with histological age estimation and be introduced to a new method that uses the anterior midshaft of the femur. In addition, newly defined and tested histological variables will be presented and distributed.

This presentation will impact the forensic science community by providing a new method for histological age estimation. This will result in a higher quality of forensic practice and reduce errors in histological analysis.

Estimating adult age is problematic owing to biological variability in skeletal age indicators and their differential response to environmental factors over an individual's life. Furthermore, when standard age indicators (e.g., pubic symphyses and sternal rib ends) are absent or altered by post-depositional taphonomic factors, anthropologists often resort to less accurate methods such as cranial suture closure. In order to improve age estimates, the use of multiple age indicators and various modalities of assessment should be considered. Histological methods are based on the continuous turnover of primary cortical bone with secondary cortical bone, which has been argued to occur at a more predictable rate than other degenerative changes. Despite this, a histological approach is typically not employed owing in part to inherent methodological issues ranging from subjective definitions to difficulty reproducing microscopic field sizes. This research evaluates histological age estimation using the anterior femur and explores the biological limitations of bone turnover as an age indicator. The study builds upon previous histological methods, recognizing their importance and impact to our current understanding of bone turnover as an age indicator.

The sample includes femur cross-sections of known age individuals from three histological collections. The sample consists of 206 individuals (102 males, 104 females) from the Ericksen collection and 14 individuals (8 males, 6 females) from the Kerley collection, and 16 (10 males, 6 females) from a modern forensic collection of known age individuals.<sup>1,2</sup> Prior to this study, research was performed to redefine and validate histological variables. The variable definitions are available to attendees as a supplemental document provided by the authors. The following variables were collected:

- Surface Area (Sa.Ar.) per mm<sup>2</sup>
- Intact Secondary Osteons (N.On.):
- Fragmentary Secondary Osteons (N.Fg.On.)
- Intact Secondary Osteon Population Density (I-OPD) per mm<sup>2</sup>
- Fragmentary Osteon Population Density (F-OPD) per mm<sup>2</sup>
- Osteon Population Density (OPD): sum of I-OPD and F-OPD
- Mean Osteonal Cross-Sectional Area (On.Ar) per mm<sup>2</sup>
- Mean Anterior Cortical Width (Ant.Ct.Wi.) per mm<sup>2</sup>

Histomorphometric data were collected using a transmitted light microscope and a firewire digital camera. The topographic sampling method was modeled after Iwaniec and colleagues and Stout and Paine.<sup>3,4</sup> The method evaluates ten columns from the periosteal to the endosteal cortex located at the anterior femur midshaft. Using a Merz counting reticule at 200x magnification (field area = 0.2304 mm<sup>2</sup>), 50% of the microscopic fields were evaluated in each column by alternating fields. This sampling strategy accounts for 95% of the remodeling variability within the anterior cross-section. Principles of stereology were followed, thus different magnifications or counting reticules may be employed to collect the histological variables. Osteon areas and cortical widths were calculated using imaging software.

Statistical analyses were performed in SPSS 19 to examine the relationship between age and cortical bone histomorphometrics. Stepwise linear regression was used to develop the prediction equation and bootstrap methods were performed to assign measures of accuracy to sample estimates. Two variables (F-OPD and On.Ar.) required log transformation to meet normality requirements. Analysis of observer error was performed using Bland and Altman's procedure for testing the repeatability of methods.<sup>5</sup>

To examine the relationship between histomorphometrics and age, a general linear model was employed. Pearson correlations show moderate and strong relationships with age for all collected variables except I-OPD. Due to this finding it was determined that the constituent variables for OPD should remain separate in the regression model. A slight level of collinearity between predictor variables was recognized and influenced the selection of variables. A one-way ANOVA indicated that all variables, with the exclusion of I-OPD ( $p=0.296$ ), demonstrate significant sex differences at the 0.05 level. Stepwise regression analysis of the male dataset produced a model using F-OPD [log] and I-OPD as predictors, while the female model selected F-OPD [log] and Ant.Ct.Wi. as predictors. The standard error of the estimate is 12.87 years and 10.49 years, respectively. In the event that sex cannot be determined, a general equation was developed using F-OPD [log] and I-OPD, providing a standard error of 11.98. Observer error results indicate the method passed repeatability standards as described by Bland and Altman.<sup>5</sup>

Current histological methods demonstrate significant issues that affect their reliability and accuracy. The method developed from this research demonstrates several advantages over previous methods. The method is based on validated



## Physical Anthropology Section - 2012

---

variable definitions, accounts for 95% of the spatial variation in osteons within the anterior cortex, and is not restricted to a specific field size or magnification. The age-related biological significance observed in the histological variables demonstrates that age-related accumulation of intact and fragmented osteons is not equivalent. While the literature suggests combining I-OPD and F-OPD to reduce observer error, the results suggest that doing so may impact the ability to interpret age-related bone remodeling. Overall, the results of the study indicate that histological analysis of the anterior femur provides reliable age estimates. Considering the biological variation in both macroscopic and microscopic adult age indicators, the standard error in this study is similar to that of previous studies with large sample sizes. One of the most prevalent issues regarding adult age estimation is the inability to accurately age older adults. The described regression model is most accurate for individuals over 50 years of age. Bearing in mind that the elderly are a rapidly growing percentage of North American populations and that unidentified adults are a common occurrence in the forensic setting, this research will significantly increase the accuracy of estimating age for older adults.

### References:

1. Erickson MF. Histological estimation of age at death using the anterior cortex of the femur. *Am J Phys Anthropol* 1991;84:171-179.
2. Kerley ER. The microscopic determination of age in human bone. *Am J Phys Anthropol* 1965; 23:149-164.
3. Iwaniec UT, Crenshaw TD, Schoeninger MJ, Stout SD, Erickson MF. Methods for improving the efficiency of estimating total osteon density in the human anterior mid-diaphyseal femur. *Am J Phys Anthropol* 1998;107:13-24.
4. Stout SD, Paine RR. Brief communication: histological age estimation using rib and clavicle. *Am J Phys Anthropol* 1992;87: 111-15.
5. Bland JM, Altman DG. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* 1986;8476:307-10.

### **Histomorphometry, Age Estimation, Forensic Anthropology**